Conference



Synthesis of Porphyrin-Peptide conjugates and interaction with $A\beta_{42}$: potential use of the systems as theranostic agents in Alzheimer's disease.

Rita Tosto,^{*a,b,**} Giuseppe Di Natale,^{*a*} Tiziana Campagna,^{*a*} Francesco Bellia,^{*a*} Giuseppe Pappalardo,^{*a*}

Alzheimer's disease (AD) is the most common cause of senile dementia affecting more than 50 million people worldwide. AD is a disorder of the central nervous system, clinically characterized by progressive loss of memory and other cognitive skills. Unfortunately, there is no cure for AD.

The major pathophysiological hallmark is the formation of amyloid deposits with a common β -sheet structure¹. Amyloid plaques are the results of a long process initiated with the seeding and production of smaller aggregated soluble forms of A β called oligomers. They are the underlying toxic species responsible for synaptic dysfunction in the brains of AD patients.

Our research focus on development of new compounds able to interfere with the early stages of the aggregation process of the A β peptides. Our compounds could be of interest as potential drug candidates AD therapy. We designed and synthesized a series of di-functional systems explicating synergic and/or additive actions to counteract the adverse effects of A β aggregated forms.^{2,3}

In particular, we covalently linked a cationic metallo-porphyrin with the well-known A β -recognizing KLVFF amino acid sequence.⁴ The peptide conjugation to the porphyrin macrocyclic was first accomplished via the formation of an amide. A second generation was obtained in good yields using a click chemistry approach.

In this communication, we describe the different synthesis approach and the ability of the conjugated porphyrin peptides to interact with $A\beta$, together with the role of metal ions (Zn or Cu) at the core of porphyrin macrocycle in assisting the recognition process.⁵

The interaction between the peptide conjugate and $A\beta$ was studied by using an array of different biophysical techniques including Dynamic Light Scattering (DLS), far-UV, Circular Dicroism (CD), Fluorescence spectroscopy and MALDI-TOF-MS.

References

- 1 W. M. Tay, D. Huang, T. L. Rosenberry, A. K. Paravastu, The alzheimer's amyloid-β(1–42) peptide forms off-pathway oligomers and fibrils that are distinguished structurally by intermolecular organization, Journal of Molecular Biology 425 (14) (2013) 2494 2508. doi:https://doi.org/10.1016/j.jmb.2013.04.003.
- 2 P. De Bona, M. Laura Giuffrida, F. Caraci, A. Copani, B. Pignataro, F. Attanasio, S. Cataldo, G. Pappalardo, E. Rizzarelli, Design and synthesis of new trehalose-conjugated pentapeptides as inhibitors of $a\beta(1-42)$ fibrillogenesis and toxicity, Journal of Peptide Science 15 (3) (2009) 220–228. doi:10.1002/psc.1109.
- 3 A. Sinopoli, A. Giuffrida, M. F. Tomasello, M. L. Giuffrida, M. Leone, F. Attanasio, F. Caraci, P. De Bona, I. Naletova, M. Saviano, A. Copani, G. Pappalardo, E. Rizzarelli, Ac-LPFFD-Th: A Trehalose-Conjugated Peptidomimetic as a Strong Suppressor of Amyloid-β Oligomer Formation and Cytotoxicity, ChemBioChem 17 (16) (2016) 1541–1549. doi:10.1002/cbic.201600243.
- 4 V. Villari, R. Tosto, G. Di Natale, A. Sinopoli, M. F. Tomasello, S. Lazzaro, N. Micali, G. Pappalardo, A Metalloporphyrin-Peptide Conjugate as an Effective Inhibitor of Amyloid-β Peptide Fibrillation and Cytotoxicity, ChemistrySelect 2 (28) (2017) 9122–9129. doi:10.1002/slct.201701148.

^a CNR - Istituto di Cristallografia, Via Paolo Gaifami 18, Catania, Italy

^b International PhD program in Chemical Sciences, University of Catania, Catania.

Creative Commons Attribuzione - Non commerciale - Condividi allo stesso modo 4.0 Internazionale

[†] oral communication at 1 st Conference on Crystallography, Structural Chemistry and Biosystems, (Catania) 04-06/10/2021

5 S. Lazzaro, N. Ogrinc, L. Lamont, G. Vecchio, G. Pappalardo, H. R. M. A., Ion mobility spectrometry combined with multivariate statistical analysis: revealing the effects of a drug candidate for Alzheimer's disease on $A\beta$ 1-40 peptide early assembly, Analytical and Bioanalytical Chemistry 411 (2019) 6353–6363. doi:10.1007/s00216-019-02030-7.